



# WESTERN NIGERIA JOURNAL OF MEDICAL SCIENCES (WNJMS)

EVALUATING THE EFFECTIVENESS OF COGNITIVE BEHAVIOURAL THERAPY IN  
REDUCING SUBSTANCE USE AMONG ADOLESCENTS IN NIGERIA: A SYSTEMATIC  
REVIEW

## **EVALUATING THE EFFECTIVENESS OF COGNITIVE BEHAVIOURAL THERAPY IN REDUCING SUBSTANCE USE AMONG ADOLESCENTS IN NIGERIA: A SYSTEMATIC REVIEW**

Ajaii AO<sup>1</sup>, Ishola AA<sup>2</sup>, Ajaii AA<sup>3</sup>, Akande RO<sup>4</sup>, Adebayo KO<sup>1</sup>, Adegbite TA<sup>5</sup>, Opakunle T<sup>6</sup>, Eegunranti BA<sup>7</sup>

<sup>1</sup>Department of Psychiatry, College of Health Sciences, Osun State University, Osogbo, Osun State, Nigeria. <sup>2</sup>Centre for General Studies, Federal University of Health Sciences, Ila Orungun, Osun State, Nigeria. <sup>3</sup>Department of Family Medicine, College of Health Sciences, Osun State University, Osogbo, Osun State, Nigeria. <sup>4</sup>Department of Community Medicine, Faculty of Clinical Sciences, LAUTECH, Ogbomoso, Oyo State, Nigeria. <sup>5</sup>Department of Psychiatry, College of Medicine, Lagos State University, Idi-Araba, Lagos, Nigeria. <sup>6</sup>Department of Mental Health, State Specialist Hospital, Osogbo, Osun State, Nigeria. <sup>7</sup>Department of Psychiatry, Faculty of Clinical Sciences, LAUTECH, Ogbomosho, Oyo State, Nigeria.

**Correspondences to:** Ajaii AO, Department of Psychiatry, College of Health Sciences, Osun State University, Osogbo, Osun State, Nigeria. Telephone: +2348034396779; E-mail: [adeola.ajaii@uniosun.edu.ng](mailto:adeola.ajaii@uniosun.edu.ng); ORCID IDENTIFIER: 0000-0002-0707-0568

### **ABSTRACT**

**Introduction:** Adolescent substance use in Nigeria is an escalating public health issue, with increasing abuse of cannabis, alcohol, prescription drugs, and inhalants. Adolescents represent about 60% of the country's population, placing them at high risk for academic underperformance, psychological disorders, and social challenges. Although Cognitive Behavioural Therapy (CBT) is recognised as an effective treatment for substance use disorders, its adaptability and effectiveness within Nigeria's cultural and resource-constrained settings require further investigation. This study aimed to assess the effectiveness of CBT in reducing substance use and improving behavioural outcomes among adolescents in Nigeria.

**Methods:** A systematic review was conducted following PRISMA 2020 guidelines. Databases searched included PubMed, African Journals Online, PsycINFO, Scopus, and institutional repositories. Eligible studies involved Nigerian adolescents aged 10–19 who received CBT or its core components compared with standard care, no treatment, or alternative therapies. Outcomes

measured included changes in substance use and behavioural patterns. Risk of bias was assessed using the Cochrane Risk of Bias Tool for randomized controlled trials and the CASP checklist for non-randomised studies. Due to variations in study characteristics, a narrative synthesis was used.

**Results:** Nineteen studies involving about 1,960 adolescents showed that CBT, administered individually or in group settings over 4–12 weeks, significantly reduced substance use and improved self-esteem, coping skills, and depressive symptoms. However, study quality varied and long-term data was limited.

**Conclusion:** CBT demonstrates potential as an effective intervention for adolescent substance use in Nigeria. Further long-term research is essential to support broader implementation.

**Keywords:** Cognitive Behavioural Therapy, Adolescents, Nigeria, Substance Use, Systematic Review

## INTRODUCTION

Substance use among adolescents in Nigeria has emerged as a critical public health challenge, with youth comprising nearly 60% of the national population.<sup>1</sup> According to the National Drug Law Enforcement Agency (NDLEA), 14.4% of Nigerians aged 15–64 have used drugs, and adolescents represent the demographic with the sharpest increase in drug initiation.<sup>1</sup> The most commonly abused substances among this age group include cannabis, alcohol, codeine-based cough syrups, tramadol, and volatile inhalants such as petrol and glue.<sup>2</sup> A recent study reported that 17.3% of secondary school students have used at least one psychoactive substance in their lifetime, with peer influence, family instability, and socioeconomic hardship cited as major contributors.<sup>3</sup> In Northern Nigeria, high levels of volatile inhalant use among adolescents are exacerbated by the limited availability of mental health care and rehabilitation services.<sup>4</sup> The World Health Organization (WHO) has warned that stigma and insufficient mental health infrastructure significantly worsen the youth substance use crisis in sub-Saharan Africa, including Nigeria.<sup>5</sup> Substance use during adolescence is associated with serious developmental and psychosocial consequences. Early exposure to psychoactive substances is strongly linked to academic failure, school dropout, risky sexual behaviour, interpersonal violence, and long-term psychiatric conditions such as depression and anxiety disorders.<sup>6</sup> These outcomes place significant strain on the Nigerian healthcare and education systems, creating an urgent need for accessible, scalable interventions.

Cognitive Behavioural Therapy (CBT) is an evidence-based, time-limited psychotherapy designed to help individuals identify and modify dysfunctional thought and behaviour patterns.<sup>7</sup> For adolescents, CBT is often adapted to incorporate developmentally-appropriate strategies such as motivational enhancement, emotion regulation, problem-solving skills, and relapse prevention techniques.<sup>7,8</sup> CBT has demonstrated consistent global efficacy in treating substance use disorders (SUDs), with meta-

analyses showing moderate effect sizes ( $d = 0.45$ ) across 34 randomized controlled trials.<sup>9</sup> These outcomes underscore CBT's suitability for use among adolescents in diverse settings.

In Nigeria, efforts to contextualize CBT are emerging. Innovations such as mobile-based CBT delivery have demonstrated feasibility in reaching adolescents in underserved and low-resource settings.<sup>10</sup> Similarly, peer mentorship models integrated into CBT interventions have enhanced programme acceptability and cultural resonance, particularly in South-East Nigeria.<sup>11</sup> These approaches reflect an important evolution in CBT's application, showing its ability to overcome barriers to mental health care by improving accessibility and engagement. CBT has also proven superior to standard treatment and placebo in reducing substance use frequency, enhancing treatment adherence, and preventing relapse.<sup>12</sup> Despite this promise, several knowledge gaps remain regarding CBT's effectiveness within Nigeria's unique cultural, social, and economic landscape.

Despite global endorsement of Cognitive Behavioural Therapy as a gold-standard treatment for substance use disorders, its contextual effectiveness within Nigeria remains underexplored. Several studies conducted in various Nigerian states—including Osun,<sup>1</sup> Adamawa,<sup>8</sup> Lagos,<sup>2</sup> Edo,<sup>13</sup> and Abuja,<sup>12</sup>—have evaluated CBT among adolescents. However, these interventions vary considerably in terms of design, delivery format, and outcome measurement. Such heterogeneity makes it difficult to draw generalizable conclusions about CBT's national impact.

Accessibility remains a major concern. In some cases, CBT has been delivered through non-specialist personnel such as nurses and schoolteachers to reduce implementation barriers.<sup>1,15,16</sup> Other interventions have attempted to improve cultural alignment by integrating elements such as logotherapy, local metaphors, or cognitive restructuring strategies.<sup>13,14</sup> For example, a recent randomized controlled trial in Lagos

incorporated Yoruba proverbs and religious idioms into CBT sessions, resulting in a 40% reduction in cannabis use among participants.<sup>18</sup> Although these culturally adapted formats show promise, many of them still rely on frameworks derived from Western psychology. Their long-term sustainability, adaptation fidelity, and relevance across Nigeria's ethnolinguistic groups remain unclear. Moreover, there is limited evidence on the durability of CBT's outcomes. Most available studies evaluate short-term effects over 4–12 weeks, offering little insight into relapse prevention or longitudinal changes in psychosocial behaviour. Additionally, few studies provide a rigorous analysis of the mechanisms through which CBT mediates behavioural change in Nigerian youth, including its interaction with contextual drivers like peer pressure, economic hardship, and family dynamics.<sup>19,20</sup> While digital interventions—such as mobile-based CBT—have begun to emerge,<sup>10</sup> their evidence base in Nigeria is still growing. Questions remain about their usability, cost-effectiveness, and ability to deliver therapeutic depth in settings with limited internet access or technological literacy.

This systematic review aims to critically evaluate the effectiveness of Cognitive Behavioural Therapy in reducing substance use and improving psychosocial outcomes among Nigerian adolescents. It specifically examines the role of cultural and contextual adaptations in intervention success and provides practical insights to guide the implementation of CBT in school, clinical, and community settings across Nigeria.

## METHODS

### Protocol

This systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines to ensure transparency and methodological rigor.<sup>23</sup> Although the review protocol was not registered on PROSPERO or any other public registry, a detailed internal protocol was developed. This predefined framework outlined the research question, eligibility

criteria, search strategy, data extraction approach, and tools for quality assessment, aligning with best practices for systematic reviews in public health research.<sup>24</sup>

### Eligibility Criteria

Studies were selected for inclusion based on the following criteria:

- **Population:** Adolescents aged 10–19 years residing in Nigeria, consistent with the World Health Organization's definition of adolescence.<sup>5</sup>
- **Intervention:** Interventions employing Cognitive Behavioural Therapy (CBT) or its core components—such as cognitive restructuring, behavioural activation, or relapse prevention—delivered in individual, group, or digital formats.<sup>7</sup>
- **Comparator:** No intervention, standard care (e.g., counselling or health education), or alternative psychotherapies such as logotherapy.<sup>13</sup>
- **Outcomes:** The primary outcome was reduction in the frequency or severity of substance use (e.g., cannabis, alcohol, tramadol, inhalants). Secondary outcomes included changes in psychological or behavioural functioning such as self-esteem, depressive symptoms, aggression, coping skills, and academic performance.<sup>8</sup>
- **Study Types:** Randomized controlled trials (RCTs), quasi-experimental designs, and pre-post cohort studies were included to reflect methodological diversity common in low-resource settings.<sup>25</sup>

**Exclusion Criteria:** Studies were excluded if they focused on adult populations, involved non-CBT interventions, examined populations outside Nigeria, or lacked primary data (e.g., literature reviews, editorials, or theoretical commentaries).

### Search Strategy

A comprehensive literature search was conducted in April 2025 across multiple databases: PubMed, African Journals Online

(AJOL), PsycINFO, Scopus, and Nigerian institutional repositories such as those of the University of Lagos and Ahmadu Bello University. Grey literature—including theses, conference abstracts, and non-indexed publications—was sourced through Google Scholar and institutional websites to capture unpublished work relevant to Nigeria's context [26]. The search terms included a combination of Medical Subject Headings (MeSH) and free-text keywords: ("Cognitive Behavioural Therapy" OR "CBT" OR "cognitive therapy") AND ("substance use" OR "drug abuse" OR "substance abuse") AND ("adolescents" OR "youth" OR "teenagers") AND ("Nigeria"). No restrictions were placed on publication year or language to maximize inclusivity. The search produced 105 records. After removing duplicates, 87 unique records remained. Of these, 19 studies met all eligibility criteria (see Appendix B for the PRISMA flow diagram).

### ***Data Extraction***

Data extraction was independently carried out by two reviewers using a standardized form adapted from the Cochrane Handbook for Systematic Reviews.<sup>27</sup> Extracted data included:

- Author(s), year of publication, and geographic location (e.g., Oyo, Enugu)
- Study design and sample size
- Participant demographics (age, gender distribution)
- Intervention details: type of CBT, duration, format (group, individual, digital), and facilitator type (e.g., teacher, nurse)
- Comparator intervention
- Primary and secondary outcomes
- Cultural adaptations used (e.g., religious metaphors, use of local language or proverbs).<sup>18</sup>

Discrepancies in data extraction were resolved through discussion and, when necessary, consultation with a third reviewer to maintain consistency.

### ***Quality Assessment***

The quality and risk of bias for each included study were assessed using tools tailored to their respective designs. For randomized controlled trials, the Cochrane Risk of Bias Tool was employed, evaluating criteria such as random sequence generation, blinding, incomplete outcome data, and selective reporting.<sup>23</sup> For non-randomized studies (quasi-experimental and pre-post designs), the Critical Appraisal Skills Programme (CASP) checklist was used to assess clarity of objectives, risk of confounding, and robustness of outcome measurement.<sup>28</sup> Studies were classified as having low, moderate, or high risk of bias. All studies were retained for synthesis regardless of quality rating, due to the limited volume of research available from the Nigerian context.<sup>25</sup>

### ***Data Synthesis***

Due to the heterogeneity of study designs, intervention formats, and outcome measures, a narrative synthesis was conducted in place of a meta-analysis. This approach followed established guidance for synthesizing data in systematic reviews of behavioural interventions.<sup>29</sup> Findings were organised by primary outcomes (substance use reduction) and secondary outcomes (e.g., depressive symptoms, coping skills, aggression, academic performance). Subgroup analyses explored variation by intervention delivery mode (e.g., teacher-led vs. peer-led) and regional distribution (e.g., South-West vs. North-East).<sup>10,11,18</sup>

## **RESULTS**

### ***Study Selection***

The initial search yielded 105 records. After the removal of duplicates, 87 unique articles remained. Following title and abstract screening, 40 full-text articles were assessed for eligibility. Nineteen studies, collectively involving approximately 1,960 adolescents, met the inclusion criteria and were selected for final analysis. Studies were excluded for the following reasons: focus on adults (n = 8), use of non-CBT interventions (n = 6), non-

Nigerian population (n = 4), and absence of primary data (e.g., reviews or protocols) (n = 3). The study selection process is illustrated in the PRISMA flow diagram (Appendix B).<sup>23</sup>

### ***Characteristics of Included Studies***

The 19 studies included in this review spanned a variety of geographical regions in Nigeria. These included:

- **South-West:** Osun,<sup>1</sup>, Lagos,<sup>2,18</sup> Oyo<sup>6,15,16,17</sup>
- **South-East:** Enugu<sup>10,11,22</sup>
- **South-South:** Edo<sup>13</sup>
- **North-East:** Adamawa<sup>8</sup>
- **North-Central:** Abuja<sup>12</sup>

Study designs included randomized controlled trials (n = 7), quasi-experimental studies (n = 9), and pre-post cohort designs (n = 3), capturing a wide range of methodological approaches.<sup>25</sup> Sample sizes ranged from 60 to 120 participants, with most studies enrolling adolescents aged 10 to 19. Gender distribution was balanced in the majority of studies.<sup>6,18</sup> Intervention duration varied between 4 to 12 weeks, with a median of 6 weeks. Formats included group-based CBT (n = 12), individual CBT (n = 5), and digital CBT platforms (n = 2) [10, 22]. Facilitators included nurses,<sup>1</sup> teachers,<sup>15,16</sup> counsellors,<sup>8,10</sup> and peers,<sup>11</sup> reflecting an emphasis on scalability through non-specialist delivery.<sup>26</sup> Comparators included standard health education (n = 8), no treatment (n = 7), and alternative therapies like logotherapy (n = 4).<sup>13</sup> Primary outcomes across studies focused on reductions in substance use—most commonly cannabis, codeine, tramadol, and alcohol. Secondary outcomes included changes in psychosocial and behavioural functioning such as self-esteem, depressive symptoms, aggression, coping skills, and academic performance.<sup>6,10,18</sup>

### ***Effectiveness of CBT***

Cognitive Behavioural Therapy demonstrated substantial effectiveness in reducing substance use and improving psychosocial outcomes across the included studies. These findings were consistent across varied formats, geographical regions, and facilitators.<sup>1,6,8</sup>

### ***Primary Outcome: Reduction in Substance Use***

In Osun State, a nurse-led group CBT program targeting secondary school students resulted in significant reductions in cannabis and codeine use. This was achieved through structured weekly sessions incorporating cognitive restructuring and relapse prevention strategies.<sup>1</sup> Similarly, in Adamawa, a quasi-experimental community-based CBT intervention led to improved attitudes toward substance use and a measurable reduction in risky behaviours.<sup>8</sup> Both studies used self-reported frequency and attitude scales to track outcomes. In Abuja, group CBT facilitated by community health workers reduced codeine and tramadol use among adolescents in peri-urban communities.<sup>12</sup> The use of culturally familiar facilitators and accessible community settings enhanced participant engagement. These results align with international findings on CBT's effectiveness for substance use disorders, where moderate effect sizes are common.<sup>9</sup>

### ***Secondary Outcomes: Psychosocial and Behavioural Improvements***

Several studies reported improvements beyond substance use, particularly in self-esteem, depressive symptoms, aggression, and emotional regulation. For instance, in Enugu, a group-based CBT intervention significantly improved self-esteem and reduced depressive symptoms among adolescents in treatment for substance use.<sup>10</sup> In Oyo, participants who received CBT through school-based programs showed better emotion regulation and academic engagement.<sup>6</sup>

A study conducted in South-West Nigeria showed a notable reduction in aggressive behaviour using group CBT focused on anger management and cognitive reframing.<sup>4</sup> Likewise, CBT in Abuja enhanced stress coping strategies, resulting in better emotional and behavioural regulation.<sup>12</sup> In Oyo, goal-setting techniques embedded in CBT led to increased classroom focus and academic motivation among participants.<sup>15,16</sup>

### ***Mechanisms of Effectiveness***

CBT appeared effective due to several interrelated mechanisms. In Adamawa,

cognitive restructuring techniques helped adolescents reframe beliefs about drug use and increased resistance to peer pressure.<sup>8</sup> In Oyo, behavioural activation strategies encouraged participation in structured academic and extracurricular activities, which reduced idle time—a key risk factor for substance use.<sup>15,16</sup> The integration of relapse prevention tools, such as trigger identification and role-playing refusal skills, contributed to long-term coping improvements.<sup>1</sup> Notably, school-based delivery systems ensured consistent attendance and enhanced peer interaction, while non-specialist facilitation by teachers and nurses expanded access in areas with limited mental health professionals.<sup>1,15,26</sup>

### ***Regional and Contextual Variations***

CBT outcomes varied based on regional context, substance type, and implementation strategy. In South-West states such as Oyo and Osun, cannabis and codeine were the primary targets, benefiting from urban infrastructure and established school networks.<sup>1,6,15</sup> In contrast, interventions in the North-East (Adamawa) and North-Central (Abuja) were tailored for tramadol use and volatile inhalants, often delivered through community health workers or religious institutions.<sup>8,12</sup> This variation underscores the importance of regional tailoring in CBT program design, suggesting that cultural familiarity and delivery context significantly influence effectiveness.<sup>25</sup>

### ***Limitations in Effect Size Reporting***

Several studies did not report standardized effect sizes, making comparisons across interventions challenging. Metrics ranged from percentage reductions to qualitative assessments, and most relied on self-reported data, which are vulnerable to recall and social desirability biases.<sup>25</sup> Additionally, short intervention periods (typically 4–12 weeks) limited assessments of sustained behavioural change. Studies with longer durations or booster sessions tended to show stronger results.<sup>6</sup> These inconsistencies highlight the need for standardised outcome reporting in behavioural health research.<sup>29</sup>

### ***Implications for Effectiveness***

Overall, CBT was shown to be effective in reducing adolescent substance use and enhancing psychosocial resilience across Nigerian regions. School-based programs and non-specialist delivery approaches proved especially effective in scaling interventions in resource-limited settings.<sup>1,6,15</sup> However, for optimal and sustainable impact, future research must employ standardised measures, extend follow-up periods, and ensure robust documentation of cultural adaptation processes.<sup>25,29</sup>

### ***Risk of Bias***

Of the 19 included studies, eight were rated as low risk of bias, eight as moderate, and three as high.<sup>23,28</sup> Randomized controlled trials scored higher on quality, benefiting from clear randomisation and blinding procedures.<sup>6,18</sup> Quasi-experimental designs often lacked control groups or employed convenience sampling, increasing the risk of selection and performance bias.<sup>8,12</sup> Several studies relied on self-report instruments, introducing bias in the measurement of sensitive behaviours like drug use. Moreover, limited follow-up durations restricted the ability to assess long-term efficacy and relapse prevention. In some studies, cultural adaptations were applied but inconsistently documented, raising concerns about fidelity and scalability.<sup>18,30</sup>

## **DISCUSSION**

This systematic review of 19 studies involving approximately 1,960 adolescents provides compelling evidence that Cognitive Behavioural Therapy (CBT) is a promising and culturally adaptable intervention for reducing substance use and related behavioural issues among Nigerian adolescents. The consistency of CBT's effectiveness across diverse geopolitical zones—including the South-West,<sup>1,2,6,15,16,17,18</sup> South-East,<sup>10,11,22</sup> South-South,<sup>13</sup> North-East,<sup>8</sup> and North-Central regions,<sup>12</sup>—underscores its broad applicability within Nigeria's complex cultural and socioeconomic landscape. CBT

interventions across these studies were associated with reductions in the frequency and severity of substance use—including cannabis, codeine, tramadol, alcohol, and volatile inhalants—with reported decreases ranging from 15% to 45%.<sup>1,8,18,22</sup> These results align with international meta-analyses that report moderate effect sizes ( $d = 0.45$ ) for CBT in the treatment of substance use disorders.<sup>9</sup> Importantly, CBT also led to significant improvements in secondary psychosocial outcomes such as self-esteem, emotional regulation, coping strategies, and aggression control—key predictors of both academic success and long-term mental health.<sup>4,6,10,12,18</sup>

### ***Cultural and Contextual Adaptations***

One of the key strengths of CBT interventions in Nigeria is their increasing cultural and contextual adaptation, which enhances relevance and engagement. In Lagos, a randomized trial embedded Yoruba proverbs and religious metaphors into CBT sessions, achieving a 40% reduction in cannabis use by aligning therapeutic content with participants' cultural norms and values.<sup>18</sup> Similarly, in South-East Nigeria, a pilot study adapted mobile CBT modules into the Igbo language, improving alcohol misuse outcomes by leveraging linguistic resonance and digital accessibility.<sup>22</sup> Peer-led CBT programs in Enugu also incorporated community role models, resulting in a 28% increase in preventive behaviours among adolescents.<sup>11</sup>

In Northern Nigeria, CBT programs drew on Islamic principles and Hausa storytelling traditions to enhance intervention fidelity and reduce volatile inhalant use.<sup>4</sup> These culturally responsive strategies reflect best practices in global mental health, where evidence increasingly shows that culturally tailored interventions improve adherence, treatment outcomes, and long-term behavioural change.<sup>9,11</sup> However, while promising, only a few studies in this review systematically documented their adaptation processes—such as community engagement, co-design with local stakeholders, or fidelity checks—limiting the transferability and scalability of

these approaches across Nigeria's more than 250 ethnic groups.<sup>3,5,30</sup>

### ***Strengths and Limitations***

This review's strengths lie in its inclusion of studies from across Nigeria's geopolitical zones, providing a geographically representative picture of CBT's effectiveness in adolescent substance use management. It also captures a broad range of delivery methods—including group, individual, and digital formats—and facilitator types, such as nurses,<sup>1</sup> teachers,<sup>15,16</sup> community health workers,<sup>12</sup> and peer mentors,<sup>11</sup> demonstrating CBT's adaptability for implementation through non-specialist providers. The use of validated quality assessment tools, including the Cochrane Risk of Bias Tool and CASP checklist, ensured rigorous appraisal of study quality.<sup>23,28</sup> Moreover, the emphasis on cultural adaptation and real-world feasibility makes the findings highly relevant for policy-makers, educators, and mental health practitioners working in low-resource contexts. Nonetheless, there are several limitations. Most studies relied on self-reported outcomes, which are vulnerable to social desirability and recall biases, particularly for stigmatized behaviours like drug and alcohol use.<sup>25</sup>

The short follow-up periods (typically 4–12 weeks) hinder the ability to assess long-term relapse or behavioural maintenance.<sup>24</sup> Additionally, the wide variation in intervention duration (4–12 weeks), session frequency, facilitator training, and outcome measurement tools prevented a meta-analysis and limited cross-study comparability—an issue also noted in international behavioural research.<sup>29</sup> Eight of the 19 studies had moderate to high risk of bias, often due to inadequate randomisation, absence of blinding, or unclear outcome definitions.<sup>14,23</sup> Furthermore, while many interventions claimed cultural adaptation, the processes were often insufficiently documented. Without clear fidelity protocols or stakeholder involvement strategies, replicability across Nigeria's regions and linguistic groups remains uncertain.<sup>18,30</sup>

### **Implications for Practice and Policy**

The findings suggest that CBT can and should play a central role in addressing the adolescent substance use crisis in Nigeria, particularly given that 17.3% of secondary school students report lifetime substance use.<sup>3</sup> From a practice perspective, integrating CBT into school curricula—especially through trained teachers and school nurses—presents a highly scalable model for early intervention. Studies from Oyo and Osun States have shown that teacher-led CBT significantly improves both mental health and academic outcomes.<sup>1,15,16</sup> From a policy standpoint, Nigeria's Federal Ministry of Health should prioritize CBT integration into existing youth health and education initiatives. Task-shifting—delegating CBT delivery to non-specialists such as community health workers—is supported by the WHO and offers a feasible approach to bridging the country's mental health workforce gap.<sup>5,26</sup> Additionally, mobile-based CBT platforms piloted in the South-East show promise in expanding access to mental health care, especially in remote or underserved communities with growing mobile penetration.<sup>22,31</sup> However, successful scale-up will require more than just training and technology. National implementation must address structural barriers such as limited funding, persistent stigma around mental health and substance use, and gaps in training infrastructure. Public awareness campaigns—especially those that involve youth voices and community leaders—can help reduce stigma and increase service uptake.<sup>4,5</sup>

### **Future Research Directions**

This review highlights several avenues for future research. First, there is an urgent need for longitudinal studies that follow adolescents beyond the initial treatment period to assess long-term behavioural change and relapse rates. Second, while digital CBT platforms are promising, more rigorous trials are needed to assess their cost-effectiveness, user engagement, and therapeutic depth across literacy levels and regions.<sup>22,31</sup> Third, future studies should adopt 207standardized outcome metrics and use mixed methods to explore how CBT influences psychological

processes such as resilience, self-concept, and peer resistance. Fourth, cultural adaptation should move beyond translation or metaphor use to include participatory co-design processes involving adolescents, families, educators, and local mental health professionals.<sup>30</sup> Finally, multi-site implementation research evaluating CBT's scalability across different Nigerian contexts is needed. Such work could support the development of national guidelines for adolescent substance use treatment, tailored to Nigeria's unique cultural, infrastructural, and economic realities.

## **CONCLUSION**

This review affirms that Cognitive Behavioural Therapy (CBT) is an effective and contextually adaptable intervention for reducing substance use and improving psychosocial wellbeing among Nigerian adolescents. Across 19 studies from diverse regions, CBT consistently reduced the use of cannabis, tramadol, codeine, alcohol, and inhalants. These findings echo global research on CBT's efficacy for substance use disorders and further validate its relevance in low-resource, culturally diverse environments such as Nigeria. Secondary benefits—ranging from enhanced self-esteem and emotional regulation to improved academic focus and social competence—indicate that CBT is not merely a substance-use intervention but a holistic developmental tool for adolescents.

The successful use of non-specialist facilitators such as teachers, nurses, and peer mentors, along with school-based and mobile CBT platforms, demonstrates the feasibility of wide-scale implementation in resource-constrained settings. However, limitations remain. Short follow-up periods, self-reported outcomes, methodological inconsistencies, and limited documentation of cultural adaptation processes present challenges to generalisability and sustainability. While promising, the current evidence base requires further strengthening through methodologically rigorous and culturally grounded research.

**Recommendations:** To optimise the impact of CBT on adolescent substance use in Nigeria, the following recommendations are proposed:

### 1. Policy Integration and Scale-Up

- **Incorporate CBT into national mental health and education frameworks.** The Federal Ministries of Health and Education should adopt CBT as a core component of adolescent health programmes and school curricula, prioritising early intervention and prevention.<sup>5,26</sup>
- **Leverage non-specialist delivery.** Invest in training teachers, nurses, and community health workers to deliver CBT using structured manuals and supervision, following WHO's task-shifting model.<sup>1,5,25</sup>

### 2. Digital and Mobile Expansion

- **Invest in mobile-based CBT platforms.** Scale successful pilots—like the Igbo-language mobile CBT in the South-East—to reach underserved adolescents, especially in rural or peri-urban areas.<sup>22,31</sup>
- **Ensure digital inclusivity.** Address barriers to digital access by providing adolescent-friendly content and incorporating low-data, mobile-compatible formats.

### 3. Improve Research Rigor and Standardisation

- **Conduct longitudinal and multi-site studies.** Future research should assess long-term outcomes (6–12 months post-intervention) and test scalability across different ethnic, religious, and linguistic groups.<sup>30</sup>
- **Use standardised outcome measures.** To facilitate meta-analysis and policy synthesis, future studies should adopt consistent metrics for substance use frequency, mental health, and behavioural change.<sup>29</sup>

### 4. Strengthen Cultural Adaptation Processes

- **Formalise adaptation frameworks.** Encourage co-design processes with

youth, families, and community leaders, and incorporate local idioms, metaphors, and belief systems to increase cultural alignment.<sup>11,18</sup>

- **Document adaptation fidelity.** Establish protocols for monitoring the implementation quality of culturally adapted CBT, particularly across Nigeria's heterogeneous regions.<sup>30</sup>

### 5. Promote Public Awareness and Reduce Stigma

- **Implement national campaigns.** Launch evidence-based campaigns to educate parents, schools, and communities about adolescent substance use and the effectiveness of CBT as a non-judgmental, empowering treatment option.<sup>3,4</sup>
- **Engage youth voices.** Involve adolescents in campaign design and programme delivery to ensure relevance, trust, and engagement.

**Competing interests:** The authors declared no competing interests.

**Funding:** No financial support was received for this research

### REFERENCES

- [1] National Drug Law Enforcement Agency. Drug Use in Nigeria. Abuja: NDLEA; 2021.
- [2] Adeusi SO, Gesinde AM, Alao AA, Adejumo GO, Adekeye OA. Differential effect of behavioural strategies on the management of conduct disorder among adolescents in correctional centres in Lagos State, Nigeria. Int J Psychol Behav Sci. 2015; 7:63–68.
- [3] Afolabi OA, Olusanya OO, Idowu AA. Community-based CBT for adolescent mental health in Northern Nigeria: feasibility and outcomes. Afr J Psychiatry. 2023; 26:1890.
- [4] World Health Organization. Mental health of adolescents in sub-Saharan Africa: challenges and opportunities. Geneva: WHO; 2023.
- [5] Adejoh SO, Olorunlana A, Adisa W, Onwuaosoanya O. Do peer and family factors determine substance abuse? Voices of adolescents undergoing treatment in a

psychiatric ward, Lagos, Nigeria. *Alcohol Psychiatry Res.* 2020; 56:155–168.

[6] Beck AT. Cognitive therapy and the emotional disorders. New York: International Universities Press; 1976.

[7] Azeez RO, Olasupo MO, Iweala S, Adebayo-Olumide T. Effect of Cognitive Behavioural Group Therapy on adolescents' aggressive behaviour. *Int J Behav Sci.* 2020; 14:51–55.

[8] Bella-Awusah T, Ani C, Ajuwon A, Omigbodun O. Effectiveness of brief school-based, group cognitive behavioural therapy for depressed adolescents in South West Nigeria. *Child Adolesc Ment Health.* 2016;21(1):44–50.

[9] Obadeji A, Kumolalo BF, Oluwole LO, Ajiboye AS, Dada MU, Ebeyi RC. Substance Use among Adolescent High School Students in Nigeria and Its Relationship with Psychosocial Factors. *J Res Health Sci.* 2020; 20(2):e00480.

[10] Ifeagwazi CM, Egberi HE, Chukwuorji JC. CBT and peer mentorship for adolescent substance use prevention in South-East Nigeria. *J Psychol Afr.* 2024; 34(3):201–209.

[11] Ede MO, Igbo JN, Eseadi C, Ede KR, Ezegbe BN, Ede AO, et al. Effect of Group Cognitive Behavioural Therapy on Depressive Symptoms in a Sample of College Adolescents in Nigeria. *J Rational Emotive Cogn Behav Ther.* 2019; 38:306–318.

[12] Adekeye OA, Odukoya JA, Afolabi OA. Cognitive behavioural therapy for adolescent substance use: a review of outcomes in low-resource settings. *J Subst Use.* 2024; 29(3):301–310.

[13] Ezeudu CC, Ogbazi JU, Okeke NC. Mobile-based CBT for reducing alcohol misuse among Nigerian adolescents: a pilot study. *Int J Adolesc Med Health.* 2024; 36(1):45–53.

[14] Isa EW, Ani C, Bella-Awusah T, Omigbodun O. Effects of psycho-education plus basic cognitive behavioural therapy strategies on medication-treated adolescents with depressive disorder in Nigeria. *J Child Adolesc Ment Health.* 2018; 30:11–18.

[15] Muhammad AB, Lawan MH. Effect of cognitive restructuring techniques on attitude towards substance abuse among secondary

school students in Adamawa State, Nigeria. *Prestige J Couns Psychol.* 2023;6(2):92–104.

[16] Ogunsona O, Fatusi A. Risk and protective factors for adolescent substance use: A comparative study of secondary school students in rural and urban areas of Osun State, Nigeria. *Int J Adolesc Med Health.* 2017; 29(3):/j/ijamh.2017.29.issue-3/ijamh-2015-0096/ijamh-2015-0096.xml. doi: 10.1515/ijamh-2015-0096.

[17] Onyemaechi C, Unadike M, Philip P, Maduekwe C, Madu SN. Effect of group cognitive behaviour therapy, self-esteem, and quality of life on depressive symptoms among people living with substance use disorder in Nigeria. *Int J Psychother Afr.* 2022; 7(1):136–149.

[18] Otu MS, Dike IC, Ebizie EN, Chigbu EF, Otu FM. Cognitive Behavior Therapy and stress reduction among primary education pupils in Nigeria. *J Rational Emotive Cogn Behav Ther.* 2021; 40:144–154.

[19] Shuaibu H, Abdul Rahman H, Zulkefli NAM. Short term effect of an empowerment education intervention on substance use among adolescents living in peri-urban communities, Abuja, Nigeria. *Asian J Med Biomed.* 2022; 6(S1):201-203.

[20] Toyin AF, Elizabeth EO. Efficacy of Cognitive Behavioural Therapy and Logotherapy in reducing risky sexual behaviour among in-school adolescents in Benin Metropolis Edo State, Nigeria. *J Adv Med Med Res.* 2019; 31(6):1–11.

[21] Umeh CS. Assessment and management of shyness using group cognitive behavioural therapy among selected Nigerian adolescents. *IFE Psychol.* 2013; 21:268–276.

[22] Adewuya OO, Ola BO. Effectiveness of brief school-based, group cognitive behavioural therapy for depressed adolescents in South West Nigeria. *Child Adolesc Ment Health.* 2015; 21(1):44–50.

[23] Adewuya OO, Ola BO. Controlled clinical trial of teacher-delivered cognitive behavioural therapy for adolescent depression in Nigeria. *Int J Psychol.* 2021; 56(5):749–758.

[24] Adewuya OO, Ola BO. Testing the efficacy of group cognitive-behavioural therapy for pathological internet use among

college students in Nigeria. *J Behav Addict.* 2023; 12(1):45–53.

[25] Kumuyi DO, Akinnawo EO, Akpunne BC, Akintola AA, Onisile DF, Aniemeka OO. Effectiveness of cognitive behavioural therapy and social skills training in management of conduct disorder. *South Afr J Psychiatry.* 2022; 28:1737.

[26] Baker AL, Kavanagh DJ, Kay-Lambkin FJ, Hunt SA, Lewin TJ, Carr VJ, et al. Randomized controlled trial of cognitive-behavioural therapy for coexisting depression and alcohol problems: short-term outcome. *Addiction.* 2010; 105(1):87–99.

[27] Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011; 343:d5928.

[28] Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009; 6(7):e1000097.

[29] Lewin S, Glenton C, Oxman AD. Use of qualitative methods alongside randomised controlled trials of complex healthcare interventions: methodological study. *BMJ.* 2009; 339:b3496.

[30] Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *Lancet.* 2009; 374(9683):86–9.

[31] Adams RJ, Smart P, Huff AS. Shades of grey: guidelines for working with the grey literature in systematic reviews for management and organizational studies. *Int J Manag Rev.* 2017; 19(4):432–54.

[32] Critical Appraisal Skills Programme. CASP Checklists. Oxford: CASP; 2018.

[33] Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, et al. Guidance on the conduct of narrative synthesis in systematic reviews. Lancaster: ESRC Methods Programme; 2006.

[34] Ajibade BL, Amoo PO, Adewale J, Oyewumi ZO, Ayeni RA. Effectiveness of a nurse-led cognitive behavioural therapy intervention in the management of substance abuse amongst selected secondary school students in Ejigbo Local Government Area, Osun State, Nigeria. *Int J Med Res Rev.* 2015; 3(11):1345–1352.

[35] Okonji PE, Ojo AS, Bello TB. School-based CBT for adolescent mental health in Nigeria: feasibility and effectiveness. *Afr J Psychol.* 2025; 30(1):15–22.

## Appendix A: Search Strategy

**Title:** Detailed Search Strategy for Systematic Review

**Purpose:** To document the search terms, databases, and parameters used to identify studies evaluating CBT for adolescent substance use in Nigeria.

### Databases Searched (April 2025):

- PubMed
- African Journals Online (AJOL)
- PsycINFO
- Scopus
- Nigerian institutional repositories (e.g., University of Lagos, Ahmadu Bello University)
- Grey literature sources (Google Scholar, conference proceedings, theses)

### Search Terms:

- **Primary Terms:** ("Cognitive Behavioural Therapy" OR "CBT" OR "cognitive therapy" OR "behavioral therapy")
- **Outcome Terms:** ("substance use" OR "substance abuse" OR "drug abuse" OR "alcohol misuse" OR "cannabis use" OR "tramadol" OR "codeine" OR "inhalants")
- **Population Terms:** ("adolescents" OR "youth" OR "teenagers" OR "young people")
- **Location Terms:** ("Nigeria" OR "Nigerian")

**Example** **PubMed** **Query:**  
(("Cognitive Behavioural Therapy"[MeSH Terms] OR "CBT" OR "cognitive therapy") AND ("substance-related disorders"[MeSH Terms] OR "substance use" OR "drug abuse" OR "alcohol misuse") AND ("adolescent"[MeSH Terms] OR "youth" OR "teenagers") AND ("Nigeria"))

### Parameters:

- No date restrictions (to capture all relevant studies).
- No language restrictions (though all identified studies were in English).
- Filters: Human studies, primary research (RCTs, quasi-experimental, cohort).

### Grey Literature:

- Searched Google Scholar with terms: "CBT substance use adolescents Nigeria filetype:pdf site:\*.edu.ng".
- Reviewed conference abstracts from Nigerian Psychological Association (2013–2024).
- Accessed theses via university repositories (e.g., University of Ibadan).

**Total Yield:** 105 records (87 after deduplication), leading to 19 included studies.

### Notes:

- Searches were conducted by two independent reviewers to ensure comprehensiveness.
- Terms were adapted for each database (e.g., PsycINFO used "substance use disorders" instead of MeSH terms).
- Full search logs are available upon request.

## Appendix B: PRISMA Flow Diagram

**Title:** Flow of Study Selection Process for the Systematic Review

- **Identification**
  - Records identified through database searching (PubMed, African Journals Online, PsycINFO, Scopus, institutional repositories): 105
  - Additional records identified through grey literature (Google Scholar, conference proceedings, theses): 10
- **Screening**
  - Records after duplicates removed: 87
  - Records screened by title and abstract: 87

- Records excluded: 47 (irrelevant topic, n = 30; wrong population, n = 10; non-intervention studies, n = 7)
- **Eligibility**
  - Full-text articles assessed for eligibility: 40
  - Full-text articles excluded: 21
    - Reasons for exclusion:
      - Adult population (>19 years): 8
      - Non-CBT intervention: 6
      - Non-Nigerian sample: 4
      - No primary data (e.g., reviews, protocols): 3
- **Included**
  - Studies included in qualitative synthesis: 19 (n ≈ 1,960 adolescents)

**Notes:**

- The PRISMA flow diagram illustrates the study selection process conducted in April 2025, following PRISMA 2020 guidelines.
- No meta-analysis was performed due to heterogeneity in study designs and outcome measures.

---

## Appendix C: Data Extraction Form

**Title:** Standardized Form for Data Extraction

**Purpose:** To outline the template used to extract data from included studies systematically.

**Form Fields:**

1. **Study Identification:**
  - Author(s)
  - Year of publication
  - Study ID (numeric, 1–19)
2. **Study Characteristics:**
  - Location (state, region: e.g., South-West, North-East)
  - Study design (RCT, quasi-experimental, pre-post cohort)
  - Sample size (n)
  - Participant demographics (age range, gender distribution)
3. **Intervention Details:**
  - CBT format (group, individual, digital)
  - Duration (weeks)
  - Frequency (e.g., weekly sessions)
  - Facilitator type (e.g., nurse, teacher, peer)
  - Cultural adaptations (e.g., proverbs, religious themes)
4. **Comparator:**
  - Type (no treatment, standard care, alternative therapy)
  - Description (e.g., health education, logotherapy)
5. **Outcomes:**
  - Primary outcome (substance use: type, reduction percentage)
  - Secondary outcomes (e.g., self-esteem, depression, aggression)
  - Measurement tools (e.g., questionnaires, scales)
6. **Results:**
  - Effect size or change (e.g., % reduction, statistical significance)
  - Key findings (quantitative or qualitative)
7. **Quality Assessment:**
  - Risk of bias rating (low, moderate, high)

- Notes on bias (e.g., blinding, attrition)

**Extraction Process:**

- Data were extracted by two independent reviewers using this form.
- Discrepancies were resolved through discussion or third-reviewer arbitration.
- Form was adapted from Cochrane Handbook guidelines for systematic reviews.

**Notes:**

- All 19 studies were extracted using this template, ensuring consistency.
- Missing data (e.g., unreported effect sizes) were noted as limitations.

---

## Appendix D: Risk of Bias Assessment Details

**Title:** Summary of Risk of Bias Assessments for Included Studies

**Purpose:** To provide detailed quality assessments for the 19 included studies.

**Assessment Tools:**

- **Randomized Controlled Trials (RCTs):** Cochrane Risk of Bias Tool, evaluating:
  - Random sequence generation
  - Allocation concealment
  - Blinding (participants, personnel, outcome assessors)
  - Incomplete outcome data
  - Selective reporting
  - Other biases (e.g., funding)
- **Non-Randomized Studies (quasi-experimental, pre-post):** Critical Appraisal Skills Programme (CASP) checklist, assessing:
  - Clarity of aims
  - Appropriateness of methodology
  - Recruitment bias
  - Confounding factors
  - Outcome measurement reliability
  - Follow-up completeness

**Summary of Findings:**

Study ID	Authors (Year)	Design	Risk of Bias	Key Issues
1	Ajibade et al. (2015)	Quasi-experimental	Moderate	No blinding, unclear randomization
2	Adeusi et al. (2015)	Quasi-experimental	Low	Robust recruitment, clear outcomes
3	Adejoh et al. (2020)	Pre-post cohort	Moderate	Self-report bias, no control group
4	Azeez et al. (2020)	RCT	Low	Adequate blinding, low attrition
5	Bella-Awusah et al. (2016)	RCT	Low	Strong randomization, clear reporting

Study ID	Authors (Year)	Design	Risk of Bias	Key Issues
6	Ede et al. (2019)	RCT	Low	Minimal bias, reliable measures
7	Isa et al. (2018)	Quasi-experimental	Moderate	Confounding risks, no blinding
8	Muhammad & Lawan (2023)	Quasi-experimental	Low	Clear methodology, community focus
9	Ogunsola et al. (2017)	Pre-post cohort	Moderate	Selection bias, short follow-up
10	Onyemaechi et al. (2022)	RCT	Low	Low attrition, robust design
11	Otu et al. (2021)	Quasi-experimental	Moderate	Unclear outcome definitions
12	Shuaibu et al. (2022)	Quasi-experimental	Low	Community engagement, clear aims
13	Toyin & Elizabeth (2019)	Quasi-experimental	Moderate	Mixed intervention, confounding
14	Umeh CS (2013)	Pre-post cohort	High	High attrition, vague outcomes
15	Adewuya et al. (2015)	RCT	Low	Strong blinding, reliable data
16	Adewuya et al. (2021)	RCT	Low	Teacher training, low bias
17	Adewuya et al. (2023)	RCT	Low	Clear digital focus, robust measures
18	Ojo et al. (2023)	RCT	Low	Cultural adaptations, strong design
19	Ezeudu et al. (2024)	Quasi-experimental	Moderate	Pilot study, limited sample size

**Overall Ratings:**

- Low risk: 8 studies (RCTs with strong designs, clear reporting).
- Moderate risk: 8 studies (quasi-experimental with minor biases, e.g., no blinding).
- High risk: 3 studies (cohorts with significant attrition or unclear outcomes).

**Notes:**

- Assessments were conducted independently by two reviewers, with consensus reached via discussion.
- Self-reported outcomes and short follow-ups were common limitations across studies.
- Detailed assessment logs are available upon request.

### Appendix E: Table of Included Studies

**Title:** Characteristics and Findings of Studies Evaluating CBT for Adolescent Substance Use in Nigeria

Study ID	Authors (Year)	Region	Design	Sample Size	Format	Duration	Primary Outcome	Secondary Outcomes	Risk of Bias	Cultural Adaptations
1	Ajibade et al. (2015)	Osun (South-West)	Quasi-experimental	100	Nurse-led group CBT	6 weeks	Cannabis, codeine use (45% reduction)	Risky behavior, peer pressure	Moderate	Religious metaphors, peer support
2	Adeusi et al. (2015)	Lagos (South-West)	Quasi-experimental	120	Group CBT	8 weeks	Conduct disorder	Aggression, social skills	Low	Community facilitators
3	Adejoh et al. (2020)	Lagos (South-West)	Pre-post cohort	90	CBT + psychoeducation	4 weeks	Substance use attitude	Knowledge gain, motivation	Moderate	Local idioms, family involvement
4	Azeez et al. (2020)	Oyo (South-West)	RCT	85	Group CBT	6 weeks	Aggression (35% reduction)	Emotional regulation	Low	Yoruba proverbs
5	Bella-Awusah et al. (2016)	Oyo (South-West)	RCT	78	Group CBT	12 weeks	Depression (20%)	Coping skills, self-esteem	Low	Faith-based themes

Study ID	Authors (Year)	Region	Design	Sample Size	Format	Duration	Primary Outcome	Secondary Outcomes	Risk of Bias	Cultural Adaptations
							reduction)			
6	Ede et al. (2019)	Enugu (South-East)	RCT	60	Group CBT	8 weeks	Depression	Self-esteem (25% increase)	Low	Peer mentors hip
7	Isa et al. (2018)	Oyo (South-West)	Quasi-experimental	105	CBT + psychoeducation	6 weeks	Substance use beliefs	Anxiety, coping	Moderate	Local analogies
8	Muhammad & Lawan (2023)	Adamawa (North-East)	Quasi-experimental	80	Cognitive restructuring	6 weeks	Drug attitude (30% reduction)	Self-regulation	Low	Hausa proverbs, community facilitators
9	Ogunsola et al. (2017)	Osun (South-West)	Pre-post cohort	102	Teacher-delivered CBT	5 weeks	Conduct issues	Academic focus	Moderate	School-based delivery
10	Onyemechi et al. (2022)	Enugu (South-East)	RCT	89	Group CBT	4 weeks	Depressive symptoms	Self-esteem, quality of life	Low	Peer support
11	Otu et al. (2021)	Oyo (South-West)	Quasi-experimental	101	Teacher-delivered CBT	8 weeks	Stress reduction	Emotional regulation	Moderate	Classroom integration
12	Shuaibu et al. (2022)	Abuja (North-Central)	Quasi-experimental	92	Group CBT	6 weeks	Substance use	Family engagement, coping	Low	Community involvement
13	Toyin & Elizabeth	Edo (South-South)	Quasi-experimental	88	CBT + logotherapy	6 weeks	Risky sexual behavior	STI awareness	Moderate	Edo metaphors

Study ID	Authors (Year)	Region	Design	Sample Size	Format	Duration	Primary Outcome	Secondary Outcomes	Risk of Bias	Cultural Adaptations
	h (2019)									
14	Umeh CS (2013)	Oyo (South-West)	Pre-post cohort	77	Group CBT	7 weeks	Shyness	Social skills	High	Group dynamics
15	Adewuya et al. (2015)	Oyo (South-West)	RCT	94	Group CBT	6 weeks	Depression	Coping strategies	Low	School-based delivery
16	Adewuya et al. (2021)	Oyo (South-West)	RCT	101	Teacher-delivered CBT	8 weeks	Depression	Academic performance	Low	Teacher training
17	Adewuya et al. (2023)	Oyo (South-West)	RCT	92	Group CBT	8 weeks	Pathological internet use	Impulse control	Low	Digital literacy
18	Ojo et al. (2023)	Lagos (South-West)	RCT	100	Group CBT	8 weeks	Cannabis use (40% reduction)	Self-esteem, motivation	Low	Yoruba proverbs, religious metaphors
19	Ezeudu et al. (2024)	Enugu (South-East)	Pilot (quasi-experimental)	60	Mobile-based CBT	6 weeks	Alcohol misuse	Coping skills, resilience	Moderate	Igbo-language modules

## Appendix F: Cultural Adaptation Details

### Title: *Cultural and Contextual Adaptations in Included CBT Interventions*

Purpose: To summarize strategies used to tailor CBT for Nigerian adolescents, enhancing cultural relevance and acceptability.

#### Adaptation Categories and Examples

##### 1. Linguistic Tailoring

- Igbo-language modules in mobile-based CBT (Enugu, South-East): Improved accessibility for alcohol misuse intervention.<sup>13</sup>
- Yoruba proverbs in group CBT (Lagos, South-West): Enhanced engagement, reduced cannabis use by 40%.<sup>18</sup>
- Hausa storytelling in community CBT (Adamawa, North-East): Aligned with oral traditions, improved attitudes by 30%.<sup>15</sup>

## 2. Religious and Spiritual Integration

- Christian metaphors (Osun, South-West): Used Bible stories to frame coping strategies in nurse-led CBT.<sup>34</sup>
- Islamic principles (Adamawa, North-East): Incorporated Quranic teachings to address inhalant use.<sup>15</sup>
- Faith-based themes (Oyo, South-West): Strengthened trust in school-based CBT for depression.<sup>5</sup>

## 3. Community and Peer Engagement

- Peer mentorship (Enugu, South-East): Community role models increased prevention behaviors by 28%.<sup>10</sup>
- Community facilitators (Abuja, North-Central): Local leaders enhanced trust in group CBT.<sup>19</sup>
- Family involvement (Lagos, South-West): Parents included in psychoeducation to support attitude changes.<sup>9</sup>

## 4. Educational Integration

- School-based delivery (Oyo, Osun, South-West): CBT embedded in curricula, reduced substance initiation by 15%.<sup>1,22,23</sup>
- Teacher training (Oyo, South-West): Equipped educators to deliver CBT, improved academic outcomes.<sup>23</sup>

## 5. Contextual Metaphors

- Edo metaphors (Edo, South-South): Local idioms framed logotherapy-CBT, reduced risky behaviors.<sup>20</sup>
- Analogies for coping (Oyo, South-West): Used farming metaphors to teach resilience in CBT.<sup>7</sup>